

## Emission-Line Galaxies from the NICMOS/HST GRISM Parallel Survey

Patrick J. McCarthy<sup>1</sup>, Lin Yan<sup>1</sup>, Wolfram Freudling<sup>2</sup>, Harry I. Teplitz<sup>3,4</sup>, Eliot M.  
Malumuth<sup>3,6</sup>

Ray J. Weymann<sup>1</sup>, Matthew A. Malkan<sup>5</sup>, Robert A. E. Fosbury<sup>2</sup>, Jonathan P. Gardner<sup>3,4</sup>

Lisa J. Storrie-Lombardi<sup>1</sup>, Rodger I. Thompson<sup>7</sup>, Robert E. Williams<sup>8</sup>, Sara R. Heap<sup>3</sup>

*Subject headings:* cosmology: observations - galaxies: distances and redshifts, evolution,  
starbursts - infrared: galaxies - techniques: spectroscopic

---

<sup>1</sup>The Observatories of the Carnegie Institution of Washington,  
813 Santa Barbara St., Pasadena, CA 91101

<sup>2</sup>Space Telescope European Coordinating Facility, Karl-Schwarzschild-Str. 2. D-85748  
Garching bei Munchen, Germany

<sup>3</sup>NASA Goddard Space Flight Center, Code 681, Greenbelt, MD 20771

<sup>4</sup>NOAO Research Associate

<sup>5</sup>Astronomy Department, University of California, Los Angeles, CA 90024-1562

<sup>6</sup>Raytheon STX Corp., Lanham, MD 20706

<sup>7</sup>Steward Observatory, University of Arizona, Tucson, AZ 85720

<sup>8</sup>Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218

## ABSTRACT

We present the first results of a survey of random fields with the slitless G141 ( $\lambda_c = 1.5\mu$ ,  $\Delta\lambda = 0.8\mu$ ) grism on NICMOS. Approximately 64 square arcminutes have been observed at intermediate and high galactic latitudes. The  $3\sigma$  limiting line and continuum fluxes in each field vary from  $7.5 \times 10^{-17}$  to  $1 \times 10^{-17}$  erg cm $^{-2}$  sec $^{-1}$  and from  $H = 20$  to 22, respectively. Our median and area weighted  $3\sigma$  limiting line fluxes within a 4 pixel aperture are nearly identical at  $4.1 \times 10^{-17}$  erg cm $^{-2}$  sec $^{-1}$  and are 60% deeper than the deepest narrow-band imaging surveys from the ground. We have identified 33 emission-line objects and derive their observed wavelengths, fluxes and equivalent widths. We argue that the most likely line identification is H $\alpha$  and that the redshift range probed is from 0.75 to 1.9. The  $2\sigma$  rest-frame equivalent width limits range from 9Å to 130Å with an average of 40Å. The survey probes an effective co-moving volume of  $10^5 h_{50}^{-3}$  Mpc $^3$  for  $q_0 = 0.5$ . Our derived co-moving number density of emission line galaxies in the range  $0.7 < z < 1.9$  is  $3.3 \times 10^{-4} h_{50}^3$  Mpc $^{-3}$ , very similar to that of the bright Lyman break objects at  $z \sim 3$ . The objects with detected emission-lines have a median F160W magnitude of 20.4 (Vega scale) and a median H $\alpha$  luminosity of  $2.7 \times 10^{42}$  erg sec $^{-1}$ . The implied star formation rates range from 1 to 324 M $_{\odot}$  yr $^{-1}$ , with an average [NII]6583,6548 corrected rate of 21 M $_{\odot}$  yr $^{-1}$  for  $H_0 = 50$ km/s/Mpc and  $q_0 = 0.5$  (34 M $_{\odot}$  yr $^{-1}$  for  $q_0 = 0.1$ ).

## 1. Introduction

Spectroscopic surveys of galaxies at  $z < 1$  have reached a reasonable level of maturity in recent years (e.g. CFRS, Lilly et al. 1996; Le Fevre et al. 1996; CNOC, Yee et al. 1996). The use of specialized selection techniques has led to a similar or greater gains at redshifts of approximately 3 and larger (Steidel et al. 1996; Dickinson 1997; Hu, Cowie, & McMahon 1998), with a few examples being identified to  $z \sim 5$  (Dey et al. 1998; Weymann et al. 1998; Spinrad et al. 1999). However, to build a coherent picture of galaxy formation and evolution we are hindered by two problems. The first is that we know little of properties of normal galaxies in the region between  $1 < z < 2$ . Secondly, most high redshift galaxies now known are selected at rest-frame UV wavelengths, where the degree of extinction is uncertain (e.g. Heckman et al. 1998). The detection of heavily dust-reddened galaxies at  $z \sim 2 - 3$  in the sub-millimeter wavelength (Smail, Ivison & Blain 1997) suggests that high redshift galaxies selected by UV colors represent only part of the population at these early epochs. At present, the estimates of the global star formation rates are uncertain as they are based primarily on UV selected samples (Pettini et al. 1997; Meurer et al. 1997).

One approach to dealing with both the scarcity of spectral signatures in the visible bandpass and the uncertainties arising from extinction is to observe in the near-IR. Until recently the available detectors and spectrometers have made spectroscopic surveys in the near-IR impractical and so most programs have been based on narrow band imaging ( $\Delta\lambda \sim 1\%$ ) to search for strong emission-line objects, primarily in the K window where  $H\alpha$  can be detected for  $1.9 < z < 2.6$ . Two narrow band near infrared imaging surveys, covering several hundred square arcminutes, were carried out with the Calar Alto 3.5m telescope (Thompson, Mannucci & Beckwith 1996; Beckwith et al. 1998; Mannucci et al. 1998). The first Calar Alto survey targeted fields containing known quasars or radio galaxies and the second the redshifts of damped  $Ly\alpha$  or metal line absorption systems. The first survey discovered one emission line galaxy at  $z = 2.43$  over an area of 276 square arcminutes to a  $3\sigma$  flux limit of  $3.4 \times 10^{-16}$  erg cm $^{-2}$  s $^{-1}$ , while the second survey found 18 emission line galaxies (16 objects with detected  $H\alpha$  and 2 with [OII]3727) over an area of 163 square arcminutes to a  $3\sigma$  flux limit of  $2 \times 10^{-16}$  erg cm $^{-2}$ s $^{-1}$ . To date the deepest narrow band near-IR imaging survey is that carried out by Teplitz, Malkan & McLean (1998) with NIRC on the Keck 10m telescope. They found 13 emission line galaxies in an 11 square arcminute area to a  $3\sigma$  flux limit of  $7 \times 10^{-17}$  erg cm $^{-2}$ s $^{-1}$ . The target fields of this survey were also

selected by matching the redshifts of quasar emission and metal absorption lines to their narrow band filters. The space densities of strong  $H\alpha$  emitting galaxies inferred from these surveys span more than 2 orders of magnitude. The large disparities in depths between the Calar Alto and Keck surveys and the strong biases introduced by selecting redshifts matched to known absorption-line clouds make comparisons between the surveys and with other measures of the average star formation rate non-trivial.

NICMOS offered a unique opportunity to address this problem with multi-object spectroscopy. The wide field camera on board NICMOS (Thompson et al. 1998) is equipped with three grisms that allow slitless spectroscopy over 0.75 square arcminutes per exposure. The extremely low background at wavelengths below  $1.9\mu$  allows for a sensitive low resolution spectroscopic survey for  $H\alpha$  and other emission-lines through volumes that are an order of magnitude larger than those available to narrow-band imaging surveys due to the large redshift range sampled by the grism.

Throughout this paper,  $H_0 = 50\text{km/s/Mpc}$  and  $q_0 = 0.5$  are adopted unless explicitly indicated otherwise.

## 2. Observations

All of the data presented here were obtained with Camera 3 on NICMOS using the G141 grism and the F160W and F110W broad-band filters. The G141 spectral element covers the wavelength range from  $1.1\mu$  to  $1.9\mu$  with a mean dispersion of  $8.1 \times 10^{-3}\mu$  per pixel. The nominal resolving power of the grism is 200. The actual resolution that is achieved is a function of the apparent image size and the camera 3 PSF. With the PSF present in the parallel observations, the realized resolution is rarely better than  $R \sim 150$ . Small variations in the PSF due to changes in the optical telescope assembly (e.g. breathing) and longer term changes in the internal structure of NICMOS introduce small variations in the maximum achievable resolution. As discussed in detail below many of the objects detected in our survey are spatially resolved and the resolution of their G141 spectra is determined by the image size.

## 2.1. Parallel observing mode and scheduling

All of the observations for this investigation were obtained with HST operating in the parallel mode. While this allowed us to collect far more data than would be possible in a single primary program, it limited our ability to plan and execute the observations in a manner that optimized the science return. Useful data for this program first became available in October 1997 when the default position of the pupil alignment mirror was moved from the camera 2 optimized position to the best reachable position for camera 3. After that time camera 3 parallels were scheduled during much of the time for which WFPC2 or STIS were the primary instruments. Most of the data in this investigation was obtained as part of a public service parallel program that was implemented and managed by the Space Telescope Science Institute. The observing algorithm was quite simple. In each full orbit one of the following exposure sequences was selected: F160W and F110W imaging with exposure times of either 512 or 382 seconds, F160W ( $2 \times 256$  sec) and G141 ( $2 \times 896$  sec), or F110W ( $2 \times 191$  seconds) and G096 ( $2 \times 896$  sec). The selection of the exposure sequence for any given orbit was nearly random, but was weighted in favor of the purely imaging sequence for much of the October 1997 to March 1998 period. After that time the balance was shifted towards spectroscopy with the grisms. All of these observations were planned and implemented by the STScI staff, with the guidance of Dr. John MacKenty.

The array was read using the STEP64 read pattern. A  $2''$  dither step was executed between each pair of direct or grism exposures. This dither step was made in the detector reference frame and was along array rows. The dither step and orientation were fixed, resulting in only two independent positions regardless of the number of exposures for any given target. Small moves executed as part of the primary observing plan introduced additional offsets for some of the longer pointings.

Approximately 10% of our data were obtained as part of a crafted guest observer (GO) parallel program that was implemented for a portion of the February 1998 to May 1998 period. The basic approach was similar to the public parallel program, except that the dithering strategy was more complex. For each pointing we began with a number of F160W images with exposure times selected to equal 15% of the expected total duration of that particular pointing. Dither moves were executed between each exposure, but with step sizes and orientations that were non-redundant. The remainder of the orbit and subsequent orbits were devoted to G141 exposures with a dither pattern matched to that of the direct

images. For a given exposure time the noise levels in these crafted observations were  $\sim 20$  to 30% lower than those in the public parallels.

## 2.2. Data processing and analysis

### 2.2.1. Two dimensional spectra

We began our treatment of the data using the calibrated output from the STScI data processing pipeline. The pipeline reduction consists of dark current removal on a read-by-read basis, linearization, bad pixel masking, and cosmic ray rejection. At this stage the data remain in the format of two dimensional, single frame images for each exposure. The principal elements of the reduction process that are accomplished by the pipeline to this point are the first order bias and dark subtraction, linearization and CR rejection. Our subsequent processing improves on the bias and dark subtraction, bad pixel masking and more complete rejection of particle events. The calibrated grism exposures have not been corrected for the flat field response and have not had the sky background removed. The flat-field response is a function of wavelength for each pixel and so depends on the location of each object, and will not be the same for the object and the sky. The flat-field correction is applied during the 1-D spectral extraction.

We constructed a median sky frame for the G141 grism by combining the calibrated exposures for approximately 200 frames, covering 80 independent positions on the sky. We rejected observations made at low galactic latitudes as well as those centered on bright objects or crowded fields. This produced a high signal-to-noise uncorrected sky frame for use in the sky subtraction for each exposure. The observed background rates vary significantly from frame-to-frame depending on the ecliptic latitude and longitude. We subtracted a scaled version of the median sky from each calibrated 2-D spectral frame. The optimal scaling for the sky was found by minimizing the RMS noise in two object-free regions of each frame. Very low-spatial frequency structure remained in the background of many frames, primarily due to the shading effect in the NICMOS arrays. We removed a low order surface from each frame. This surface was derived from a second order polynomial fit to an intermediate image constructed from the median of each column. This surface fitting

was carried out separately in each quadrant of the detector, allowing us to remove small quadrant-to-quadrant bias variations. The individual sky subtracted and shading corrected frames were then shifted into registration, using offsets derived from the accompanying direct images. The registered 2D spectra were then averaged with a bad pixel mask and  $3\sigma$  clipping applied to remove hot pixels and residual cosmic rays.

The direct images that are used to calibrated the extracted one dimensional spectra were processed and combined using the same techniques as above, but without the sky subtraction and low order surface fitting correction steps.

### *2.2.2. 1-D spectral extraction and calibration*

The extraction and calibration of the one-dimensional spectra was carried out with the NICMOSlook software developed at ST-ECF (Freudling et al. 1998). The extraction was done in the following way. First, the position and size of the objects previously identified as spectral line objects were interactively marked on the image display. The extraction region of the spectrum was then computed using transformations derived as part of the calibration program. The extracted spectra were corrected for the wavelength dependent flat-field response on a pixel-by-pixel basis. For each pixel, the quantum efficiency for the extracted wavelength was estimated by interpolation from a series of narrow-band flat field exposures. Fluxes were calibrated by applying throughput curves previously derived from the observation of flux calibrators. The final one dimensional spectra were obtained by adding the flux of all pixel which correspond to each wavelength bin. For some objects close to one of the edges of the grism images, only partial spectra could be extracted. Since there is a significant displacement between the location of objects on the direct images and their dispersed spectrum on the grism image, spectra for a few objects that fall outside the area covered by the direct images were extracted. In that case, the wavelength scale was estimated from the cut-off of the throughput curves at high and low wavelengths which can easily be identified. For such objects, the wavelength scale is uncertain by about  $0.01\mu$ .

The rms accuracy of the wavelength calibration for G141 is about  $2 \times 10^{-3}\mu$ . Repeated calibration exposures have shown no changes larger than 3% in the wavelength zero point or 1.5% in the dispersion. The flux calibration is based on observations of both white dwarfs and solar analogs during the NICMOS camera 3 campaigns. The accuracy of the

spectrophotometry is limited by variations in the quantum efficiency within individual pixels. We estimate that the flux scale for an individual spectrum is accurate to about 10%.

Spectra were also extracted in a parallel effort by members of the team at the NASA Goddard Space Flight Center (GSFC) using software independent of the NICMOS-Look routines. The GSFC software followed a similar strategy to NICMOS-Look. Spectral positions on the chip were determined by the known offsets from the object location in the direct image. Extractions were performed on individual (unregistered) frames first, with registration left to the end. Each spectrum was extracted in a weighted box, similar to the procedure for the NICMOS-Look point source extraction. The wavelength solution was confirmed by independent analysis of the same calibration data. Flat-fielding was performed using the same data-cube of narrow-band observations, interpolated to the wavelengths of the extracted pixels, and the flux calibration was performed based on independent analysis of the white dwarf observations. Finally, spectra were combined in wavelength space. Emission-line wavelengths derived from spectra extracted with the GSFC software were found to agree with the NICMOSLook results to within one pixel ( $0.008\mu\text{m}$ ). Line fluxes and equivalent widths were measured using Gaussian fits to the line and polynomial fitting of the underlying continuum. The overall results of the extraction and analysis using the GSFC and ECF/NICMOSLook software were similar.

### 3. Results

#### 3.1. The properties of the data and the identification of emission-lines

There are several aspects of the data which negatively impact our ability to identify genuine spectral features. Since we are observing in the slitless mode, each object produces zero, first and second order spectra. The first order spectra contain the useful data, the zero and second orders are sources of confusion, the zero order images as they can be mistaken for emission features, particularly when they fall on the first order continua of other objects. The zero order images are slightly dispersed and for point sources often appear bimodal. The displacement between the zero order images and the start of the first order spectrum is  $27''$ , or 135 pixels, a substantial fraction of the array size. For this



reason the zero order images are confined to roughly half of the detector area and in most cases they can be identified by matching them with either the first order spectra or with an object in the direct image. There is a small portion of the detector in which zero order images can appear without either first order spectra or images in the direct frame. In addition to the confusion caused by the zero order images and second order spectra, there are artifacts associated with hysteresis in the HgCdTe detectors. Bright stars imaged with the broad-band filters just before a grism observation will produce a low level after-image that is displaced from the first order spectrum. Since the data are nearly always taken in pairs of direct and grism exposures, these persistent images appear in multiple exposures and can take on the appearance of isolated emission-lines. Strong cosmic rays events have similar properties, but the two position dithering sequence aids in their identification and removal. Particularly strong particle events effect more than one pixel and can produce amorphous clouds of electrons that decay over time scales of an orbit. During the South Atlantic Anomaly (SAA) passage the particle event rate is quite large and most NICMOS images or spectra obtained at this time are not usable. The hysteresis in the detectors often renders subsequent exposures useless as well. For a significant number of our short pointings one of the two 896 second grism exposures has greatly enhanced noise from residual CR events associated with the SAA passage.

Given the wide variety of point-like features that appear in the two dimensional grism images we have made our emission-line identifications on the basis of visual inspection. The eye can detect compact features to quite low signal to noise levels. To convince ourselves that any possible emission line was real we proceed to eliminate all possible false signals, zero order images, persistent images, CRs etc. To facilitate the detection of incompletely rejected cosmic rays and other noise features we split each observation into two frames of equal length and compared the position of possible emission features. Those features that appeared in only one of the data sets, or did not move with the dither pattern, were rejected. In Figure 1 we show two pairs of F160W/G141 frames and mark examples of each of the sources of confusion as well as real emission features.

### 3.2. Areal coverage and depth

During the period from October 1997 to October 1998, 3500 camera 3 exposures were taken as part of the public and our crafted GO parallel programs. These contain a mix of direct exposures with the F160W and F110W filters and spectroscopic exposures with the G141 and G096 grisms. A total of 456 pairs of F160W/G141 frames were taken during this time. These were distributed in 107 independent pointings with exposure times ranging from 1790 seconds to 21,480 seconds. The G141 grism spectra can provide useful data for our purposes to quite low galactic latitudes. Rather than adopt a strict latitude cutoff we chose to reduce all of the data and reject those with high stellar densities. After some experience we rejected all fields with stellar densities greater than 50 per square arcminute above a  $10\sigma$  threshold in the F160W direct image. We are left with 85 pointings, covering 64 square arcminutes.

The final depth achieved varied considerably from field to field. For the 1790 second exposures the noise level varied by a factor 3. Some of this variation was due to genuine background variations, but much of it was due to various non-Poisson sources of noise (e.g. SAA passage). For longer pointings the depth increased roughly as the square root of the exposure time. We define our limiting depths as  $3\sigma$  within a 4 pixel (0.16 square arcseconds) aperture. This area is a reasonable representation of the area of the camera 3 PSF at the focus achieved outside of the campaign mode. In Figure 2 we show a histogram of the  $3\sigma$  4-pixel limits for the G141 data.

### 3.3. Detected Emission Features

In Table 1 we list all of the fields for which we have reduced and examined the G141 grism data. For each pointing we give the J2000 IAU designation, the galactic latitude and the exposure times for the F160W and G141 filters.

In Figures 1a and 1b we show two typical pairs of direct and 2-D grism images. These two grism images show many of the features discussed in the previous section (zero order images, first and second order spectra, after images) as well as one emission-line object each. After careful examination of all of the spectra we assembled a catalog of 33 emission-line detections. Table 2 gives the field names, object designation, J2000 coordinates, observed




Fig. 1.— Two typical pairs of direct and 2-D grism images. Each field is  $51'' \times 51''$  and oriented in arbitrary directions. In the G141 frames we have marked zero order images (Z), residual after images (AI), persistent cosmic rays (CR), first and second order spectra and our identified emission-lines. The region marked as “confusion” in the upper panel identifies the portion of the detector within which zero-order images neither have corresponding first-order spectra within the G141 field of view or direct images in the F160W field. We have marked the objects producing the emission lines in the F160W images.

wavelengths, redshifts, line fluxes, significance levels, observed equivalent widths and equivalent width limits ( $2\sigma$ ) for these 33 emission line galaxies. We set a rejection threshold at  $3\sigma$ , the derived significance levels of our cataloged emission-lines are given in Table 2. These 33 emission-lines are distributed in 22 fields, 6 of which contain more than one emission-line object. We do not convincingly detect more than one line in any object. All of lines are quite strong, in terms of equivalent width. This is expected given the low spectral resolution of the grisms; weak lines against bright continua are not easily detected. The observed equivalent width limits reflect the signal-to-noise ratios in the continua as well as the galaxy sizes. The  $2\sigma$  observed equivalent width limits are calculated as follows:  $W_{lim} = 2 \times (\frac{s}{snr}) \times \sqrt{N}$ , with  $s$  the scale in  $\text{\AA}$  per pixel,  $snr$  the signal-to-noise ratio per pixel of the continuum at the emission line and  $N$  the number of pixels under each line. For lines that are not velocity resolved, the apparent line width is the convolution of the instrumental spatial resolution and the intrinsic size of the emission line region. For our emission line galaxies, we take  $N$  to be 3 pixels. The  $2\sigma$  rest-frame equivalent width limits range from  $9\text{\AA}$  to  $130\text{\AA}$  with an average of  $40\text{\AA}$ . In Figure 3, we show direct images in the F160W band ( $\sim$  H band) and 2D grism spectra for all 33 emission line galaxies. Figure 4 presents the extracted one dimensional spectra of all 33 emission line galaxies.




Fig. 2.— Histogram of the achieved  $3\sigma$  4-pixel noise levels in the G141 data. Each pointing covers 0.75 square arcminutes.



Fig. 3.— A mosaic of F160W band direct images and 2D grism spectra of 33 emission line galaxies found in our survey. The emission-lines in each object are marked with vertical ticks. For a few of the objects the continuum images fall near the edge or just off the edge of the F160W field of view. For these objects (0917+8142C, 0931-0449A) we do not have F160W images.




Fig. 3.— continued.




Fig. 4.— The extracted 1-D spectra of the emission line galaxies. For each object we have marked our candidate emission-lines with arrows below the line. We plot the entire range of the G141 grism for each spectrum even though parts of the spectrum have fallen beyond the field of view of the detector for several objects. The redshifts, assuming an H $\alpha$  identification for the line are given for each object. There are deficiencies in the background removal for several objects that lead to significant uncertainties in the continuum level (e.g. 1237+6215A)




Fig. 4.— continued.






Fig. 4.— continued.

We believe that most, if not all, of the emission lines that we have detected are either  $H\alpha$  or an unresolved blend of  $H\alpha + [\text{NII}]\lambda 6583/6548$ . In Table 3 we list the line luminosities, star formation rates derived from the Kennicutt (1984; 1992) relation, the apparent H-band magnitudes (on the Vega scale) and J–H colors for the fields where the F110W observations were available. The identification of the emission-lines as  $H\alpha$  is based on a number of arguments. First, the most plausible identifications are  $H\alpha + [\text{NII}]\lambda 6583/6548$ ,  $[\text{OIII}]\lambda 5007, 4959$ ,  $H\beta\lambda 4861$  and  $[\text{OII}]\lambda 3727$ . At the peak of the G141 transmission these correspond to redshifts of 1.28, 2.0, 2.1 and 3.0, respectively. Studies of nearby galaxies show that  $H\alpha + [\text{NII}]\lambda 6583/6548$  is usually the strongest line, the second being  $[\text{OII}]\lambda 3727$  (Kennicutt 1992). The equivalent width of  $H\beta$  scales with the equivalent width of  $H\alpha$  as  $W(H\beta) = 0.15W(H\alpha) - 4$ . Even in the bluest Magellanic irregulars and starburst galaxies,  $W(H\beta)$  is only  $\sim 5\text{\AA}$ . With a resolution of  $5000 \text{ km sec}^{-1}$ , it would be quite difficult to detect such lines in our data. Redshifts of  $\sim 3$  are highly unlikely given the median F160W magnitude of 20.5, as shown in Figure 5. The  $z \sim 3$  Lyman break galaxies have typical F160W magnitudes that are 2 magnitudes fainter ( $H \sim 22.5$ ) (Giavalisco et al. 1998). The space density of our emission-line objects is sufficiently large that they are unlikely to represent the extreme tail of the luminosity distribution of the  $z \sim 3$  Lyman break objects. If not  $H\alpha$ , it is more likely that the lines that we see are  $[\text{OIII}]\lambda 5007, 4959$  rather than  $[\text{OII}]\lambda 3727$ . The  $[\text{OIII}]\lambda 5007, 4959$  lines could be marginally resolved in our spectra, but only if the objects are spatially compact. At  $z \sim 1.5$ , if we take an  $R - H$  color of 3, the median H magnitude of our objects corresponds to  $M_R = -21.7$  for  $H_0 = 50 \text{ km/s/Mpc}$ ,  $q_0 = 0.5$ , quite close to  $M_R^*$  determined from the Las Campanas Redshift Survey (Lin et al. 1996). In Figure 6, the left panel shows the distribution of  $H\alpha$  line-luminosity; the median line-luminosity is  $2.5 \times 10^{42} \text{ erg s}^{-1}$ , close to the value of  $L_{H\alpha}^*$  derived in local surveys (e.g. Gallego et al. 1995, Tresse & Maddox 1998). If the emission lines are actually  $[\text{OIII}]\lambda 5007, 4959$ , the implied rest-frame H-band luminosities and emission-line luminosities would be a factor of 2.5–3 larger and the likelihood of such objects appearing in flux limited samples is suppressed by the exponential cutoff in the luminosity function. Strong  $[\text{OIII}]\lambda 5007, 4959$  emission is more often identified with AGN than star-forming galaxies, making it even less likely that  $[\text{OIII}]\lambda 5007, 4959$  emitters at  $z \sim 2$  dominate our detections (see section 4). Furthermore, in the redshift range between 1.6 and 1.8, both  $[\text{OIII}]\lambda 5007, 4959$  and  $[\text{OII}]\lambda 3727$  are visible in the G141 passband. Lines with longer rest-wavelengths are possible, but unlikely identifications.  $\text{HeII}\lambda 10830$  implies redshifts that are implausibly low ( $\sim 0.38$ ) and for most redshifts both of the  $[\text{SIII}]\lambda 9069, 9545$  lines would be within the passband. Teplitz et al. (1999) have confirmed the identification of  $H\alpha$  in

two of our objects (J0055+8518a,  $z = 0.76$ , J0622-0018a,  $z = 1.12$ ) by detecting [OII]3727 at the appropriate redshifts in spectra taken with LRIS on the Keck 10m telescope.

H $\alpha$  is blended with [NII]6583,6548 in our spectra. To correct for the contribution from the [NII] doublet, we adopted [NII]6583/H $\alpha$  = 0.3, the average ratio for all types of galaxies in the Gallego et al. (1997) objective-prism survey. Taking [NII]6583/[NII]6548 = 3, we have  $f(\text{H}\alpha) = 0.71 \times f((\text{H}\alpha + \text{N}[\text{II}]))$ . Thus the correction to the estimated star formation rates in Table 3 is 29%. We have not applied any extinction corrections to either the H $\alpha$  luminosities or the star formation rates.

### 3.4. Notes on Individual Fields

In Figure 3 we present cut-out images and 2-D spectra of all of our emission-line candidates. Flux and wavelength calibrated 1-D spectra are shown for the same objects in Figure 4. We designate each object by its IAU J2000 field designation and a letter. There are a few fields that have noteworthy features and we detail these below.

0040+8505 A: This is one of the weaker lines, in terms of  $W_\lambda$ , in our sample. There are only two exposures of this field, but the emission feature is seen in both frames.

0613+4752 A: The emission-line in this object is more compact than the continuum. As in the case of 0040+8505 there are only two exposures for this field, but the narrow emission feature is seen in both frames and moves with the dither pattern.

0622-0018 A: This low redshift object is detected in a very low latitude field ( $b = -6^\circ$ ). The emission-line is spatially extended and appears to follow the continuum distribution as revealed in the F160W image.

0738+0507 A: This is also a low latitude field and the object in question is the brightest in our sample. The large equivalent width and compact morphology of the emission-line suggests that this object might be an AGN.

0738+0507 B: The putative emission-line is quite faint and diffuse. There are 6 exposures of this field and the emission feature appears with equal strength in the two independent subsets of the data.

0741+6515 A: The emission-line in this object is spatially resolved and its morphology is




Fig. 5.— The distribution of apparent H magnitude of emission line galaxies. The magnitudes are on the Johnson system.




Fig. 6.— The distributions of redshifts and  $H\alpha$  line luminosities of the emission line galaxies.

similar to that of the continuum shown in the F160W image.

0741+6515 B: The emission-line in this object is at the extreme red end of the continuum spectrum and is spatially offset from the continuum. The emission-feature is located in a portion of the field such that it cannot be a zero-order artifact.

0917+8142 A: This emission feature is also at the extreme red end of the continuum. At the location of this feature there is the possibility of zero order images. If this feature were a zero order image, the object would be just off the edge of the F160W direct image. The first order continuum of a zero order image at this location would be partially on the G141 frame, but there is a confusing signal from a bright galaxy at this location. The spatial coincidence between the continuum and emission features led us to classify this as a real emission-line, although this object should be considered less secure than the others.

0917+8142 C: The putative emission-line in this object is at the edge of the G141 field of view. The feature is only present in one of the two dithered frames, it was moved off the detector by the 2'' dither motion. The object is also located just outside the F160W field in both dither positions. The great strength of this line makes it a candidate AGN.

1120+2323 A,B: The two emission-lines in this field are extremely compact. Both emission features are seen in the two dither positions. For reasons that are not entirely clear, the focus in this particular set of exposures was unusually good. The difference may have arisen from breathing of the telescope optical assembly, or from motions within the NICMOS dewar.

1143-8036 A,B: This pair of emission-line galaxies appear to have nearly identical redshifts in the 2-D G141 frame. The slightly different locations in the F160W image result in different wavelength scales, and hence different redshifts. The close proximity of the two images make accurate centroiding of the F160W images difficult and this introduces uncertainties into the wavelength calibrations.

## 4. Discussion

This survey provides a unique window on galaxies at intermediate redshifts. In so doing it provides us with a view of the bright end of the  $H\alpha$  luminosity function averaged

over a cosmologically significant volume. From these data we will derive an  $H\alpha$  luminosity function and a measure of the volume averaged star formation rate that is fairly insensitive to the effects of extinction. These analyses require careful treatment of the selection efficiencies and biases inherent in a slitless low-resolution survey and are presented in Yan et al. (1999). In the present work we restrict ourselves to a first-order census of bright emission-line objects and a brief comparison with other surveys in this redshift range.

There are a number of near-IR emission-line imaging surveys that are either in progress or have been recently completed. In Table 4 we compare the characteristics of our survey with several ground-based programs. Clean comparisons are difficult since our survey is of a different character and spans a very wide range of exposure times from field to field. In our deepest pointings we achieve flux limits that are substantially deeper than the narrow-band imaging survey being carried out by Teplitz et al. (1998) with the Keck Telescope. Our median and area weighted sensitivities are both  $4.1 \times 10^{-17}$  erg cm $^{-2}$  sec $^{-1}$  ( $3\sigma$ , 4 pixels),  $\sim 60\%$  deeper than the area-weighted detection limit reported by Teplitz et al. (1998). The two principal differences between our survey and those listed in Table 4 are the volumes probed and the choice of sight-lines. Because our survey is spectroscopic we are able to probe large columns in the  $z$  direction. The comparable areal coverage of our survey then leads to total co-moving volumes that are one to two orders of magnitude larger than those surveyed by the narrow-band imaging programs.

Inspection of Table 4 reveals that most of the narrow-band imaging surveys are targeted at lines of sight and redshifts that are *a priori* known to contain either luminous galaxies or absorption line systems. The motivation for this can be clearly seen in the results of the first Calar Alto survey (Thompson et al. 1996). Surveys along sight-lines without absorbers, or in completely random areas, yield few strong line emitters in the volumes that can be probed by imaging with 1% filters. Thompson et al. detected a single object in 30,000  $h_{50}^{-3}$ Mpc $^3$ , implying a space density of  $3 \times 10^{-5}$   $h_{50}^3$ Mpc $^{-3}$ . The choice of AGN and/or absorber redshifts is effective in producing detections, but it clearly impacts the inferred space density of line emitters. Teplitz et al. (1998), targeting a mix of QSO and absorber redshifts, report the highest space density, at approximately  $10^{-2}$   $h_{50}^3$ Mpc $^{-3}$ . This is quite different from the results of the Thompson et al. (1996) survey, even when allowing for the very different depths of the two surveys. The second Calar Alto survey (Mannucci et al. 1998) demonstrates the effect of targeting metal line absorption systems, while Bechtold et al. (1997) have achieved similar results using damped Ly $\alpha$  redshifts.

Our H $\alpha$  survey probes an effective co-moving volume of  $1.0 \times 10^5 h_{50}^{-3} \text{ Mpc}^3$  ( $2.1 \times 10^5 h_{50}^{-3} \text{ Mpc}^3$  for  $q_0 = 0.1$ ). Thus the co-moving number density of emission-line galaxies averaged over our volume is  $3.3 \times 10^{-4} h_{50}^3 \text{ Mpc}^3$ . In comparison, the present-day co-moving number density of galaxies brighter than  $L^*$  in the B band is  $7 \times 10^{-4} h_{50}^3 \text{ Mpc}^{-3}$  (Ellis et al. 1996) and the bright Lyman break galaxies at  $z \sim 3$  reported by Steidel et al. (1996) have a space density of  $3.6 \times 10^{-4} h_{50}^3 \text{ Mpc}^{-3}$ . The duty cycle of strong H $\alpha$  emission in galaxies with bursting or variable star formation histories can be quite low, given the sensitivity of strong line emission to only the most recent star formation. Thus it is not unreasonable that our space density is below that of present day  $L^*$  galaxies and comparable to that of the UV continuum selected galaxies at high redshift. The low resolution of the NICMOS grisms imprints a strong selection effect against emission-lines with low equivalent widths. Given the incompleteness at low equivalent widths and the rather bright H-band magnitudes of the detected emission-line objects it seems likely that we are sampling star formation in a population that is closely related to the majority population of massive galaxies.

The average star formation rate in our sample is  $30 M_{\odot} \text{ yr}^{-1}$  ( $40 M_{\odot} \text{ yr}^{-1}$  for  $(q_0 = 0.1)$ ). After correcting for [NII] contamination, this rate is reduced to  $21 M_{\odot} \text{ yr}^{-1}$ . We have not applied any extinction correction to the calculations of H $\alpha$  luminosities and star formation rates. While our sample is not magnitude limited, the distribution of apparent magnitudes is sufficiently narrow to allow a reasonable comparison with magnitude limited samples. As noted above, our median F160W magnitude of 20.5 is quite close to  $M_R^*$  as derived locally. It is interesting to note that the average star formation rate in the Lyman break objects, before corrections for extinction are  $\sim 8 M_{\odot} \text{ yr}^{-1}$ , and that the extinction corrected rates are a factor of 3 – 7 higher. Recent near-infrared spectroscopy of 8 CFRS galaxies at  $z \sim 1$  by Glazebrook et al. (1998) reached an H $\alpha$  luminosity of  $2 \times 10^{42} \text{ erg/s/cm}^2$  with a star formation rate of  $22 M_{\odot} \text{ yr}^{-1}$ , comparable to what we find, although our survey covers a wider range of H $\alpha$  luminosities and star formation rates.

One source of uncertainty in using our data to derive global star formations rates is the unknown degree of contamination from AGNs. The fraction of AGNs in our sample could, in principle, be calculated using the AGN luminosity function in the redshift range from 0.7 to 1.9. However, at  $z = 1.5$  an H magnitude of 20.5 corresponds roughly to an absolute B magnitude of -22.5 adopting a B–H color similar that for QSOs observed by Neugebauer et al. (1986). At this faint magnitude, the quasar luminosity function is completely unknown.

The most recent quasar luminosity function at  $z \sim 0.5 - 2.0$  reaches only to  $M_B \sim -25$  (Hewett, Foltz & Chaffee 1993). Furthermore, our knowledge of the Seyfert luminosity function is entirely based on local samples and the degree of luminosity evolution in the Seyfert galaxy populations as a function of redshift is unknown. A crude estimate of the fraction of AGNs in faint redshift surveys can be derived from spectrophotometry of the CFRS sample. Using various emission line ratios, Hammer et al. (1997) and Tresse et al. (1996) estimated that about 8–17% of their galaxies at  $z < 0.7$  are active galaxies, either Seyfert 2 or LINERs. A contamination rate at this level is consistent with our classification of two of our objects as candidate AGNs on the basis of their large equivalent widths and compact morphologies. If we exclude the two AGN candidates, the average SFR is  $28.5 \text{ M}_\odot/\text{yr}$  and  $20 \text{ M}_\odot/\text{yr}$  if we correct the  $[\text{NII}] \text{ } 6583, 6548 \text{ \AA}$  contamination. The impact of line misidentifications is comparable to that of the AGNs, as the most likely contaminant are  $[\text{OIII}] \text{ } 5007, 4959$  emission from compact non-stellar sources.

## 5. Summary

We have presented the basic data derived from a spectroscopic survey of field galaxies in the near infrared. The G141 grism on NICMOS provides a unique opportunity to survey large volumes to faint flux levels in  $\text{H}\alpha$  and other emission-lines. Our survey of 64 square arcminutes at high and intermediate galactic latitudes resulted in the detection of 33 emission-line objects and we believe that most or all of these are  $\text{H}\alpha$  at redshifts between 0.75 and 1.9. The line luminosities, equivalent widths and continuum magnitudes suggest that we are seeing active star formation in galaxies from the upper end of the luminosity function. The apparent space density of  $\text{H}\alpha$  emitters in our survey is similar to that of the bright Lyman break objects identified at  $z > 3$  and is within a factor of  $\sim 2$  of that of present day  $L^*$  galaxies. The grisms on NICMOS provide a first look at the enormous potential of low resolution near-IR spectroscopy from space.



## 6. Acknowledgments

We thank the staff of the Space Telescope Science Institute for their efforts in making this parallel program possible. In particular we thank Peg Stanley, Doug van Orsow, and the staff of the PRESTO division. We also thank John Mackenty and members of the STScI NICMOS group for crafting the exposure sequences. We also acknowledge the role of Duccio Macchetto and the Parallel Working Group, lead by Jay Frogel, in making the public parallel program a success. This research was supported, in part, by grants from the Space Telescope Science Institute, GO-7498.01-96A and P423101. HIT and JPG acknowledge funding by the Space Telescope Imaging Spectrograph Instrument Definition Team through the National Optical Astronomy Observatories and by the NASA Goddard Space Flight Center.

## REFERENCES

- Bechtold, J., Elston, R., Yee, H. K. C., Ellingson, E., & Cutri, R. 1998, in *The Young Universe*, ed. S. D’Odorico, A. Fontana, & E. Giallongo, ASP Conference Series, 146, 241
- Beckwith, S. V. W., Thompson, D., Mannucci, F., Djorgovski, S. G. 1998, *ApJ*, 504, 107
- Dey, A., Spinrad, H., Stern, D., Graham, J.R., Chaffee, F. 1998, *ApJ*, 498, 93L
- Dickinson, M. 1998, in *The Hubble Deep Field*, eds. M. Livio, M. Fall, and P. Madau, STScI, in press
- Freudling, W. & Pirzkal, N. 1998, in *NICMOS and the VLT*, eds. Wolfram Freudling and Richard Hook, p47
- Ellis, R., Colless, M. Broadhurst, T. Heyl, J. & Glazebrook, K., 1996, *MNRAS*, 280, 235
- Gallego, J., Zamorano, J., Rego, M., Vitores, A.G., 1997, *ApJ*, 475, 502
- Giavalisco, M., et al. 1998, in preparation.
- Glazebrook, K., Blake, C., Economou, F., Lilly, S. & Colless, M. 1998, astro-ph/9808276
- Hammer, F., Flores, H., Lilly, S.J., Crampton, D., Le Fevre, O., Rola, C., Mallen-Ornelas, G., Shade, D. & Tresse, L. 1997, *ApJ*, 481, 49
- Heckman, T., Robert, C., Leitherer, C., Garnett, D.R., van der Rydt, F. 1998, *ApJ*, 503, 646
- Hewett, P., Foltz, C.B. & Chaffee, F.H. 1993, *ApJ*, 406, L43
- Hu, E.M., Cowie, L.L. & McMahon, R.G. 1998, *ApJ*, 502, L99
- Kennicutt, R.C., Bothun, G.D. & Schommer, R.A. 1984, *AJ*, 89, 1279
- Kennicutt, R.C. 1992, *ApJ*, 388, 310
- Le Fevre, O., Hudson, D., Lilly, S.J., Crampton, D., Hammer, E. & Tresse, L. 1996. *ApJ*, 461, 534
- Lilly, S.J., Le Fevre, O., Hammer, F. & Crampton, D. 1996, *ApJ*, 460, L1

- Lin, H., Kirshner, R.P., Shectman, S.A., Landy, S.D., Oemler, A., Tucker, D.L., Schechter, P.L. 1996, 464, 60
- Malkan, M. A., Teplitz, H., McLean, I. S. 1996, ApJ, 468, 9
- Mannucci, F., Thompson, D.J., Beckwith, S.V.W., Williger, G.M., 1998, ApJ, 501, 11
- Meurer, G.R., Heckman, T.M., Lehnert, M.D., Leitherer, C. & Lowenthal, J. 1997, AJ, 114, 54
- Pettini, M., Kellogg, M., Steidel, C.C., Dickinson, M., Adelberger, K.L., Giavalisco, M. 1998, ApJ, in press
- Smail, I., Ivison, R.J., Blain, A.W. & Kneib, J.P. 1998, ApJL, 507, L21
- Spinrad, H., Stern, D., Bunker, A., Dey, A., Lanzetta, K., Yahil, A., Pascarelle, S. & Fernández-Soto, A. 1999, AJ, in press
- Steidel, C., Giavalisco, M., Pettini, M., Dickinson, M., Adelberger, K. 1996, ApJ, 462, L17
- Teplitz, H. I., Malkan, M. A., McLean, I., S. 1998, ApJ, in press
- Teplitz, H. I., McLean, I., S., & Malkan, M. A. 1999a, ApJ, submitted
- Teplitz, H. I. Malkan, M. A., & McLean, I. S. 1999b, in “After The Dark Ages”, in press
- Thompson, D.J., Mannucci, F. & Beckwith, S.V.W. 1996, AJ, 112, 1794
- Thompson, R.I., Rieke, M., Schneider, G., Hines, D.C., Corbin, M.R. 1998, ApJ, 49
- Tresse, L., Rola, C., Hammer, F., Stasinska, G., Le Fevre, O., Lilly, S.J. & Crampton, D. 1996, MNRAS, 281, 847
- Weymann, R., Stern, D., Bunker, A., Spinrad, H., Chaffee, F., Thompson, R.I., Storrie-Lombardi, L.J. 1998, ApJ, 505, L95
- van der Werf, P.P., Bremer, M. N., Moorwood, A. F. M., Rottgering, H. J. A. 1997, IAU, 186, 197
- Yan, L., McCarthy, P. J., Freudling, W., Teplitz, H. I., Malumuth, E. M., Weymann, R. J., Malkan, M. A., 1999, submitted to ApJ, Letters.

Yee, H.K.C., Ellingson, E. & Carlberg, R.G. 1996, ApJS, 102, 269

---

This manuscript was prepared with the AAS L<sup>A</sup>T<sub>E</sub>X macros v4.0.

TABLE 1  
THE NICMOS GRISM SURVEY FIELDS

| Field      | b  | F160W | G141  | Field      | b  | F160W | G141  | Field      | b  | F160W | G141  | Field      | b  | F160W | G141  |
|------------|----|-------|-------|------------|----|-------|-------|------------|----|-------|-------|------------|----|-------|-------|
| J0002–1531 | 73 | 2550  | 5370  | J0535–0519 | 19 | 510   | 1790  | J1120+2328 | 69 | 510   | 1790  | J1604+4318 | 48 | 3830  | 5370  |
| J0017+1544 | 46 | 1020  | 1790  | J0535–0623 | 19 | 5100  | 14320 | J1120+2324 | 69 | 3570  | 12530 | J1612+6539 | 40 | 510   | 1790  |
| J0022–7202 | 44 | 510   | 1790  | J0540–6939 | 31 | 2550  | 3580  | J1120+1300 | 64 | 16860 | 10740 | J1633–6732 | 13 | 1530  | 5370  |
| J0023–7209 | 44 | 1530  | 1790  | J0543–6909 | 31 | 1530  | 1790  | J1129+5832 | 55 | 510   | 1790  | J1633–0941 | 24 | 2550  | 8950  |
| J0029+8516 | 22 | 1020  | 1790  | J0547–7004 | 30 | 510   | 1790  | J1133+0403 | 60 | 510   | 1790  | J1642+3944 | 41 | 510   | 1790  |
| J0040+8457 | 22 | 1530  | 1790  | J0548–5101 | 30 | 510   | 1790  | J1134+0406 | 60 | 510   | 1790  | J1701–2143 | 12 | 1020  | 3580  |
| J0040+8505 | 22 | 510   | 1790  | J0551+1552 | 05 | 2550  | 3580  | J1143–8036 | 18 | 1000  | 1917  | J1726–4821 | 07 | 2338  | 6136  |
| J0050–5207 | 65 | 2040  | 1790  | J0551+1549 | 05 | 510   | 1790  | J1145+4733 | 65 | 2550  | 8950  | J1741–0701 | 12 | 510   | 1790  |
| J0051+8523 | 22 | 510   | 1790  | J0609+2415 | 02 | 2040  | 1790  | J1145+4735 | 65 | 1510  | 1917  | J1744–2900 | 00 | 510   | 1790  |
| J0055+8518 | 22 | 3060  | 10740 | J0613+4752 | 13 | 510   | 1790  | J1146+4723 | 66 | 1530  | 3580  | J1752–1739 | 04 | 2040  | 1790  |
| J0056+8510 | 22 | 510   | 1790  | J0622–0018 | 06 | 1530  | 5370  | J1151+5509 | 59 | 1031  | 2428  | J1753–1739 | 04 | 1020  | 3580  |
| J0100–7213 | 44 | 1020  | 3580  | J0627–6512 | 26 | 1020  | 1790  | J1153+4917 | 65 | 510   | 1790  | J1755–2158 | 01 | 510   | 1790  |
| J0100–7212 | 44 | 2295  | 8055  | J0655+7910 | 26 | 2040  | 7160  | J1218+4714 | 68 | 3570  | 7160  | J1755–2200 | 01 | 510   | 1790  |
| J0100–7213 | 44 | 1785  | 6265  | J0709+4840 | 22 | 510   | 1790  | J1226+3335 | 81 | 510   | 1790  | J1758+6715 | 29 | 1530  | 3580  |
| J0107+3230 | 30 | 510   | 1790  | J0738+0507 | 12 | 1530  | 5370  | J1228+4410 | 72 | 510   | 1790  | J1802–2330 | 00 | 510   | 1790  |
| J0133+3028 | 31 | 510   | 1790  | J0741+6515 | 29 | 1031  | 2428  | J1236+6225 | 54 | 510   | 1790  | J1822+1511 | 13 | 1020  | 3580  |
| J0133+3026 | 31 | 510   | 1790  | J0804+0510 | 18 | 1020  | 1790  | J1237+6219 | 54 | 6120  | 21480 | J1822+1512 | 13 | 1020  | 3580  |
| J0138–1749 | 75 | 510   | 1790  | J0808+7459 | 31 | 1510  | 1917  | J1237+6215 | 54 | 4590  | 16110 | J1822+1512 | 13 | 510   | 1790  |
| J0143+3259 | 28 | 1020  | 3580  | J0809+7608 | 30 | 510   | 1790  | J1237+6210 | 54 | 1530  | 5370  | J1824–3020 | 08 | 2040  | 3580  |
| J0233–0231 | 55 | 510   | 1790  | J0811+7505 | 31 | 510   | 1790  | J1238–4954 | 12 | 5100  | 10740 | J1847–0451 | 01 | 510   | 1790  |
| J0234+0340 | 50 | 510   | 1790  | J0851+1151 | 32 | 1660  | 1790  | J1238–4954 | 12 | 2040  | 7160  | J1900–3637 | 17 | 1020  | 1790  |
| J0234+0350 | 50 | 1020  | 3580  | J0911–6457 | 11 | 510   | 1790  | J1238–4954 | 12 | 1530  | 5370  | J1912–5948 | 25 | 1031  | 2428  |
| J0238+1634 | 39 | 2550  | 8950  | J0917+8142 | 31 | 510   | 1790  | J1256+2204 | 84 | 1020  | 1790  | J1917–3330 | 19 | 1020  | 3580  |
| J0240–0141 | 53 | 1020  | 3580  | J0920+6909 | 38 | 510   | 1790  | J1301+5900 | 58 | 510   | 1790  | J1938–1621 | 17 | 510   | 1790  |
| J0240–0140 | 53 | 1530  | 5370  | J0922+8136 | 32 | 510   | 1790  | J1316+2911 | 84 | 510   | 1790  | J1958+4039 | 05 | 5100  | 14320 |
| J0318–6632 | 44 | 510   | 1790  | J0923+8137 | 32 | 510   | 1790  | J1326–4258 | 19 | 510   | 1790  | J1958+4037 | 05 | 4080  | 10740 |
| J0325–3618 | 56 | 1020  | 3580  | J0923+8149 | 32 | 510   | 1790  | J1326–4245 | 19 | 510   | 1790  | J2000+2236 | 03 | 510   | 1790  |
| J0349+1712 | 28 | 1530  | 1790  | J0927–0502 | 31 | 1530  | 5370  | J1336+5150 | 63 | 1530  | 3580  | J2044–3124 | 36 | 2040  | 5370  |
| J0354+0945 | 32 | 1020  | 3580  | J0931–0450 | 32 | 510   | 1790  | J1336–2959 | 31 | 1020  | 3580  | J2044+6758 | 15 | 255   | 895   |
| J0400–4433 | 48 | 1020  | 3580  | J0931–0449 | 32 | 510   | 1790  | J1337+7023 | 46 | 2040  | 1790  | J2044+6758 | 15 | 1275  | 895   |
| J0408–1207 | 41 | 510   | 1790  | J0950+7222 | 38 | 510   | 1790  | J1338+7009 | 46 | 510   | 1790  | J2056–0443 | 29 | 1530  | 1790  |
| J0408–1209 | 41 | 510   | 1790  | J0954+6905 | 40 | 1318  | 2556  | J1341+2824 | 78 | 4080  | 14320 | J2106–2411 | 39 | 2040  | 3580  |
| J0421+1926 | 21 | 1530  | 1790  | J0955+6908 | 40 | 1510  | 1917  | J1343+0218 | 62 | 510   | 1790  | J2144+2813 | 18 | 510   | 1790  |
| J0424+2707 | 15 | 510   | 1790  | J0956+6937 | 40 | 1530  | 3580  | J1343+0124 | 61 | 1020  | 3580  | J2150+2845 | 19 | 1530  | 5370  |
| J0425+1709 | 21 | 510   | 1790  | J1020+3654 | 56 | 1020  | 3580  | J1343+0225 | 62 | 510   | 1790  | J2151+2854 | 19 | 1020  | 3580  |
| J0430+2449 | 15 | 510   | 1790  | J1039+4313 | 58 | 510   | 1790  | J1345+0201 | 61 | 1020  | 3580  | J2155+6334 | 07 | 1530  | 1790  |
| J0431+1803 | 20 | 1530  | 3580  | J1039+4145 | 59 | 6380  | 3580  | J1346+0206 | 61 | 510   | 1790  | J2155+6341 | 07 | 1530  | 3580  |
| J0432+2424 | 15 | 1020  | 1790  | J1044–5936 | 00 | 1530  | 1790  | J1347+0219 | 61 | 1020  | 3580  | J2155+6343 | 07 | 1530  | 3580  |
| J0433+2526 | 15 | 510   | 1790  | J1044–5948 | 00 | 3060  | 7160  | J1402+5419 | 59 | 510   | 1790  | J2156+6345 | 07 | 2040  | 1790  |
| J0443+3612 | 06 | 1530  | 3580  | J1053–0342 | 48 | 1530  | 5370  | J1415+1910 | 69 | 2040  | 5370  | J2157+6331 | 06 | 1318  | 2556  |
| J0455+3029 | 08 | 510   | 1790  | J1055–0418 | 48 | 1020  | 1790  | J1432+0423 | 56 | 1530  | 5370  | J2157+6338 | 07 | 7140  | 5370  |
| J0458–0404 | 26 | 510   | 1790  | J1056–0337 | 48 | 1020  | 3580  | J1454+1218 | 57 | 510   | 1790  | J2217–0336 | 46 | 510   | 1790  |
| J0506+5250 | 07 | 2040  | 3580  | J1057–0318 | 48 | 2040  | 5370  | J1504+0110 | 48 | 510   | 1790  | J2240+8031 | 19 | 510   | 1790  |
| J0526–6602 | 33 | 2550  | 3580  | J1120+2327 | 69 | 510   | 1790  | J1507+0234 | 49 | 1530  | 5370  | J2257+2050 | 34 | 510   | 1790  |
|            |    |       |       |            |    |       |       |            |    |       |       | J2319+0809 | 48 | 1020  | 3580  |

TABLE 2  
EMISSION-LINE OBJECTS FROM THE NICMOS GRISM SURVEY

| Field      | Obj | $\alpha(2000)$ |    | $\delta(2000)$ |     | $\lambda$<br>$\mu$ | $z_{H\alpha}$ | flux <sup>a</sup><br>erg/s/cm <sup>2</sup> | $N_\sigma$ | $W_{obs}$<br>Å | $W_{lim}^b$<br>Å |      |     |
|------------|-----|----------------|----|----------------|-----|--------------------|---------------|--------------------------------------------|------------|----------------|------------------|------|-----|
| J0040+8505 | a   | 00             | 40 | 43.93          | 85  | 05                 | 49.4          | 1.727                                      | 1.63       | 1.5            | 4.8              | 250  | 62  |
| J0055+8518 | a   | 00             | 55 | 37.85          | 85  | 18                 | 17.3          | 1.160                                      | 0.76       | 5.3            | 5.2              | 240  | 25  |
| J0240-0140 | a   | 02             | 40 | 10.85          | -01 | 40                 | 53.0          | 1.839                                      | 1.80       | 0.7            | 12               | 160  | 49  |
| J0613+4752 | a   | 06             | 12 | 59.93          | 47  | 52                 | 43.3          | 1.341                                      | 1.04       | 1.3            | 5.8              | 60   | 60  |
| J0622-0018 | a   | 06             | 22 | 13.78          | -00 | 18                 | 25.5          | 1.402                                      | 1.14       | 2.9            | 4.0              | 403  | 55  |
| J0627-6512 | a   | 06             | 27 | 32.20          | -65 | 12                 | 35.0          | 1.742                                      | 1.65       | 1.8            | 3.5              | 176  | 86  |
| J0738+0507 | a   | 07             | 38 | 57.86          | 05  | 07                 | 35.2          | 1.210                                      | 0.84       | 16             | 10.9             | 161  | 32  |
| J0741+6515 | b   | 07             | 38 | 58.37          | 05  | 07                 | 56.7          | 1.77                                       | 1.69       | 0.9            | 3.0              | 180  | 105 |
|            | a   | 07             | 41 | 40.89          | 65  | 15                 | 23.8          | 1.612                                      | 1.45       | 1.0            | 7.4              | 92   | 72  |
|            | b   | 07             | 41 | 45.37          | 65  | 15                 | 36.2          | 1.879                                      | 1.86       | 1.8            | 12.8             | 165  | 118 |
| J0917+8142 | c   | 07             | 41 | 44.67          | 65  | 15                 | 06.9          | 1.33                                       | 1.06       | 1.8            | 4.1              | 879  | 101 |
|            | a   | 09             | 17 | 47.25          | 81  | 42                 | 21.4          | 1.880                                      | 1.86       | 2.9            | 9.4              | 375  | 131 |
|            | b   | 09             | 17 | 44.96          | 81  | 42                 | 40.6          | 1.883                                      | 1.87       | 0.9            | 4.4              | 224  | 87  |
| J0923+8149 | c   | 09             | 17 | 39.64          | 81  | 43                 | 06.5          | 1.364                                      | 1.07       | 56             | 115              | 737  | 32  |
|            | a   | 09             | 23 | 34.32          | 81  | 50                 | 11.2          | 1.572                                      | 1.39       | 4.2            | 7.0              | 547  | 114 |
|            | a   | 09             | 31 | 25.97          | -04 | 49                 | 45.4          | 1.299                                      | 0.98       | 24             | 14.4             | 550  | 49  |
| J0931-0449 | a   | 10             | 39 | 41.79          | 41  | 45                 | 04.4          | 1.636                                      | 1.49       | 1.8            | 10.7             | 410  | 118 |
| J1039+4145 | a   | 10             | 39 | 41.79          | 41  | 45                 | 04.4          | 1.636                                      | 1.49       | 1.8            | 10.7             | 410  | 118 |
| J1056-0337 | a   | 10             | 56 | 56.41          | -03 | 37                 | 03.9          | 1.126                                      | 0.72       | 4.6            | 3.1              | 168  | 36  |
| J1120+1300 | a   | 11             | 20 | 37.36          | 13  | 00                 | 07.1          | 1.376                                      | 1.10       | 0.2            | 6.8              | 2809 | 60  |
| J1120+2323 | a   | 11             | 20 | 24.29          | 23  | 22                 | 40.8          | 1.557                                      | 1.37       | 1.9            | 4.8              | 660  | 74  |
|            | b   | 11             | 20 | 22.36          | 23  | 23                 | 13.0          | 1.397                                      | 1.13       | 1.4            | 4.8              | 456  | 83  |
| J1134+0406 | a   | 11             | 34 | 04.71          | 04  | 06                 | 33.7          | 1.260                                      | 0.92       | 5.4            | 3.2              | 1165 | 86  |
| J1143-8036 | a   | 11             | 43 | 19.56          | -80 | 36                 | 46.2          | 1.538                                      | 1.35       | 1.2            | 5.5              | 310  | 80  |
|            | b   | 11             | 43 | 19.73          | -80 | 36                 | 46.9          | 1.549                                      | 1.36       | 0.7            | 5.4              | 330  | 53  |
|            | c   | 11             | 43 | 10.31          | -80 | 37                 | 06.9          | 1.668                                      | 1.54       | 1.2            | 3.8              | 353  | 73  |
|            | d   | 11             | 43 | 18.46          | -80 | 36                 | 30.2          | 1.49                                       | 1.27       | 1.3            | 4.1              | 377  | 298 |
| J1145+4733 | a   | 11             | 45 | 29.14          | 47  | 33                 | 35.2          | 1.257                                      | 0.92       | 1.1            | 2.5              | 250  | 184 |
| J1218+4714 | a   | 12             | 18 | 36.42          | 47  | 14                 | 34.0          | 1.56                                       | 1.38       | 0.7            | 3.9              | 108  | 36  |
| J1237+6209 | a   | 12             | 37 | 49.04          | 62  | 09                 | 50.0          | 1.802                                      | 1.74       | 0.9            | 5.2              | 172  | 51  |
| J1237+6215 | a   | 12             | 37 | 38.85          | 62  | 15                 | 24.8          | 1.778                                      | 1.71       | 0.4            | 3.2              | 108  | 24  |
| J1237+6219 | a   | 12             | 37 | 06.57          | 62  | 19                 | 37.1          | 1.557                                      | 1.37       | 1.8            | 6.0              | 607  | 65  |
|            | b   | 12             | 37 | 05.28          | 62  | 19                 | 38.5          | 1.348                                      | 1.05       | 1.3            | 2.8              | 546  | 164 |
|            | c   | 12             | 37 | 07.71          | 62  | 19                 | 24.8          | 1.734                                      | 1.64       | 0.4            | 4.2              | 580  | 303 |

<sup>a</sup>The H $\alpha$  line flux is in unites of  $10^{-16}$  ergs/s/cm<sup>2</sup>.

<sup>b</sup> $W_{lim}$  is the  $2\sigma$  observed equivalent width limit, which reflects the signal-to-noise ratio of the continuum near the wavelength of H $\alpha$  line.

TABLE 3  
PHOTOMETRY OF EMISSION-LINE OBJECTS

| Field      | Obj | $z_{H\alpha}$ | H<br>mag | J–H<br>mag | $\log_{10}(L_{H\alpha})$<br>erg s <sup>−1</sup> | SFR<br>M <sub>⊙</sub> /yr |
|------------|-----|---------------|----------|------------|-------------------------------------------------|---------------------------|
| J0040+8505 | a   | 1.63          | 20.4     | ...        | 42.40                                           | 22                        |
| J0055+8518 | a   | 0.76          | 19.7     | ...        | 42.20                                           | 14                        |
| J0240−0140 | a   | 1.80          | 20.8     | 1.1        | 42.16                                           | 13                        |
| J0613+4752 | a   | 1.04          | 18.1     | ...        | 41.90                                           | 7                         |
| J0622−0018 | a   | 1.14          | 19.1     | ...        | 42.34                                           | 19                        |
| J0627−6512 | a   | 1.65          | 20.4     | 0.8        | 42.49                                           | 27                        |
| J0738+0507 | a   | 0.84          | 17.9     | ...        | 42.78                                           | 54                        |
|            | b   | 1.69          | 22.0     | ...        | 42.21                                           | 14                        |
| J0741+6515 | a   | 1.45          | 21.1     | 0.8        | 42.10                                           | 11                        |
|            | b   | 1.86          | 22.3     | 0.7        | 42.60                                           | 36                        |
|            | c   | 1.06          | 19.6     | 1.0        | 42.06                                           | 10                        |
| J0917+8142 | a   | 1.86          | 20.2     | ...        | 42.81                                           | 58                        |
|            | b   | 1.87          | 20.2     | ...        | 42.31                                           | 18                        |
|            | c   | 1.07          | ...      | ...        | 43.56                                           | 324                       |
| J0923+8149 | a   | 1.39          | 19.1     | ...        | 42.69                                           | 44                        |
| J0931−0449 | a   | 0.98          | 19.0     | 0.7        | 43.11                                           | 115                       |
| J1039+4145 | a   | 1.49          | 20.2     | 1.4        | 42.39                                           | 22                        |
| J1056−0337 | a   | 0.72          | 18.8     | 1.1        | 42.10                                           | 11                        |
| J1120+1300 | a   | 1.10          | 20.8     | 0.6        | 41.14                                           | 1                         |
| J1120+2323 | a   | 1.37          | 20.5     | ...        | 42.33                                           | 19                        |
|            | b   | 1.13          | 21.1     | ...        | 42.01                                           | 9                         |
| J1134+0406 | a   | 0.92          | 19.9     | 1.0        | 42.40                                           | 22                        |
| J1143−8036 | a   | 1.35          | 21.4     | ...        | 42.11                                           | 12                        |
|            | b   | 1.36          | 20.5     | ...        | 41.89                                           | 7                         |
|            | c   | 1.54          | 20.8     | ...        | 42.24                                           | 16                        |
|            | d   | 1.27          | 21.0     | ...        | 42.09                                           | 11                        |
| J1145+4733 | a   | 0.92          | ...      | ...        | 41.71                                           | 5                         |
| J1218+4714 | a   | 1.38          | 20.3     | 0.9        | 41.90                                           | 7                         |
| J1237+6209 | a   | 1.74          | 23.3     | 0.5        | 42.24                                           | 15                        |
| J1237+6215 | a   | 1.71          | 20.8     | 1.0        | 41.87                                           | 7                         |
| J1237+6219 | a   | 1.37          | 20.5     | ...        | 42.31                                           | 18                        |
|            | b   | 1.05          | 21.9     | ...        | 41.91                                           | 7                         |
|            | c   | 1.64          | 22.3     | ...        | 41.83                                           | 6                         |

NOTE.—The H $\alpha$  luminosities and star formation rates were calculated for  $H_0 = 50\text{km/s/Mpc}$  and  $q_0 = 0.5$ . Here the listed star formation rates have not been corrected for the contamination from [NII]6584,6548 line. As described in detail in the text, the correction to the star formation rates is 29%. We only have J–H colors for some of the fields where the F110W observations were available. “...” indicates no measurements.

TABLE 4  
EMISSION-LINE SURVEYS IN THE NEAR-IR

| SITE             | Type    | $\delta\lambda$ | Area<br>$\square'$ | Volume<br>Mpc <sup>3</sup> | $z$ range    | $f_{lim}^a$<br>ergs/s/cm <sup>2</sup> | $L_{lim}^b$<br>ergs/s | N <sup>c</sup> | Ref | Remarks                    |
|------------------|---------|-----------------|--------------------|----------------------------|--------------|---------------------------------------|-----------------------|----------------|-----|----------------------------|
| Calar Alto # 1   | Imaging | 1%              | 276                | 30,000                     | 1.5 - 2.8    | 3.4                                   | 3.8 – 19              | 1              | 1   | Targeted toward AGN        |
| Calar Alto # 2   | Imaging | 1%              | 228                | 20,600                     | 0.9 - 2.4    | 2.4                                   | 1.0 – 9.5             | 18             | 2   | QSO metal line absorbers   |
| Keck/NIRC        | Imaging | 1%              | 12                 | 815                        | 2.3 - 2.5    | 0.7                                   | 2.5 – 3.2             | 13             | 3   | QSO emission and absorbers |
| ESO near-IR      | Imaging | 1%              | 48                 | 10,000                     | 2.2 - 2.3    | 1.9                                   | 6.2 – 6.8             | 2              | 4   | Radio Galaxies             |
| Lick/UCLA-camera | Imaging | 1%              | 180                | 4255                       | 0.95 - 0.995 | 3.9                                   | 1.8 – 3.3             | 2              | 5   | R.Q. QSO emission z        |
| NICMOS Parallel  | GRISM   | 45%             | 58                 | 100,000                    | 0.8 - 1.9    | 0.4                                   | 0.1 – 0.9             | 33             | 6   | Random fields              |

<sup>a</sup>All the flux limits are  $3\sigma$  and in unites of  $10^{-16}$  ergs/s/cm<sup>2</sup>.

<sup>b</sup>The luminosity limits represent the minimum line luminosities which the survyes are sensitive to and are in unites of  $10^{42}$ ergs/s.

<sup>c</sup>Number of emission line galaxies found in the survey.

NOTE.—Co-moving volumes are calculated for  $H_0 = 50$  km/s/Mpc and  $q_0 = 0.5$ .

REFERENCES.—(1) Thompson et al. 1996; (2) Mannucci et al. 1998; (3) Teplitz et al. 1998; (4) van der Werf et al. 1997; (5) Teplitz et al. 1999; (6) This paper